

INPOP10a

by A. Fienga^{1,2}, H. Manche¹, P. Kuchynka¹, J. Laskar¹, M. Gastineau¹.

September 21, 2010

¹ Astronomie et Systèmes Dynamiques, IMCCE-CNRS UMR8028, Paris, France

² Observatoire de Besançon, UTINAM-CNRS UMR6213, Besançon, France

The INPOP10a version has several improvements in the fitting process, the data sets used in the fit and in the general features of the solution. No big change was brought in the dynamics. As a consequence of these changes, the extrapolation capabilities of INPOP10a are improved compared to INPOP08, especially for the Earth, Mars, Mercury and Saturn orbits. As for INPOP08, INPOP10a provides to the user, positions, velocities of the planets and the moon and TT-TDB chebychev polynomials at <http://www.imcce.fr/inpop>.

1 The Data sets

Several data sets have been added since INPOP08. The global distribution of the data used for the INPOP fit has changed its balance compared to INPOP06: now, more than 56% of the planetary data used in the fit are deduced from tracking data of spacecrafts including range deduced from tracking, VLBI angular positions and flyby normal points. The statistics of the obtained postfit residuals are presented in table 1 and 9. Comments are given in section 4. Postfit residuals are plotted in figures 3, 4, 5, 6, and 9,

1.1 Mercury

- Two normal points deduced from the Mariner tracking data in 1974 and 1975 have been provided by JPL (Folkner 2010). These points have been included in the JPL DE solutions but not distributed. They bring constraints only in range with an accuracy of about 150 meters. This is an improvement of a factor 5 compared to the previous direct radar range observations of the Earth-Mercury distances.
- Based on planetary ephemerides refitted by the Messenger navigation team to the tracking data of the Messenger mission during its flybys of Mercury (NAVPE004, NAVPE005 and NAVPE006 as provided by the SPICE NAIF), we extracted 3 normal points giving 3 very accurate geocentric positions of Mercury. These positions are given in geocentric $(\alpha, \delta, \text{range})$ with an accuracy of about 1.5 mas, 2.5 mas and 5 meters respectively.

These 5 points change drastically our knowledge of the Mercury orbit. Until now, only direct radar ranging on the Mercury surface were available with an accuracy of about 800 meters.

1.2 Mars and Venus

Like with INPOP08, tracking data of MEX and VEX missions are provided by ESA (Morley 2009, Morley 2010) based on a procedure described in Fienga et al. (2009). For extrapolation tests, we decided to keep apart the last 6 months of MEX and VEX data in order to test the extrapolation capabilities of INPOP10a (see section 4.2).

1.3 Saturn

To the Cassini normal points provided by JPL over the 2005 to 2007 period and used in INPOP08, are added VLBI observations of the spacecraft published by (Jones et al. 2010). These new data are spread over a complementary period of time: 2 points before 2005 and 8 positions obtained from 2007 to 2009.5. They provide very accurate ICRF geocentric (α, δ) with accuracies better than 1 mas.

1.4 Jupiter, Uranus and Neptune

Flybys data of the 3 planets obtained during several missions (Pioneer 10 and 11, Viking 1 and 2, Ulysses and Cassini) are also added, provided by Folkner (2009). The mean accuracy for the Jupiter flybys in range is about 2 kilometers and the data are spread over 25 years. The range flybys for Uranus and Neptune are unique points with an accuracy of about several meters. These observations improve the estimations of the geocentric distances to the outer planets while no observation of that type were used in INPOP06 and INPOP08 adjustments. New optical data obtained from 2000 to 2008 by the United States Naval Observatory Flagstaff Station using the Flagstaff Astrometric Scanning Transit Telescope are also added for Uranus, Neptune and Pluto.

1.5 Pluto

Stellar occultations are taken into account in INPOP10a by the use of measured offsets in topocentric (α, δ) . These offsets have been provided by Sicardy (2009). The accuracy of these observations is about 45 mas for α and 20 mas for δ . Such accuracy is at least a factor 2 better than the modern CCD observations.

1.6 Moon

Lunar Laser Ranging observations from 1969 to 2010 are taken into account, provided by several sites: Cerga (Grasse, France), Mc Donald, MLRS1, MLRS2 (Fort Davis, Texas, USA), Haleakala (Hawaii, USA) and Apollo (New Mexico, USA). Their accuracy reaches today a few centimeters. Since INPOP08, observations from Apollo and Cerga have been added. The weights of recent Cerga's observations have been split into 2 periods (before and after 1995), as for MLRS1 ones (before and after 1985).

2 The fitting procedure

2.1 Fit of the mass of the sun for a fixed AU

In INPOP10a, the value of the AU is fixed to the value given in the IERS2003 convention. The GM of the sun is fitted to the observations with the initial conditions of planets, the densities of the asteroids and the oblateness coefficient J_2 of the sun. Values of the fitted parameters are given in table 3.

2.2 Fit of a new selection of asteroids

The selection of asteroids modeled in INPOP10a is based on Kuchynka et al. (2010). The perturbations of 24635 asteroids have been taken into account using an averaging of their perturbations by a ring with fixed physical characteristics for most of them and individual perturbations for 161 of them (see Kuchynka et al. 2010). Based on a study of the correlations between the asteroids, we found 30 asteroids among the most perturbing objects highly correlated with each others. In order to decrease the uncertainties on the mass estimations, we fixed 15 asteroid masses to values well determined by other methods and found in Baer (2010). Besides these fixed values, we fit the 146 GMs of asteroids using a BVLS algorithm (Lawson and Hanson, 1995) with large constraints on the densities.

3 General features

3.1 IAU2009 Current Best Estimates (CBE)

We used the mass of planets provided by the IAU 2009 CBE lists (Luzum 2010).

3.2 Fit of parameters related to LLR observations

More than 180 parameters have a significant effect on LLR computations, including initial conditions, potential coefficients, reflectors and stations coordinates, biases,... But because of large correlations or better estimations obtained by other techniques, only 59 of them have been fitted. They have been chosen so that the ratio between the formal error and the fitted value is less than 5%. Values of parameters and formal errors are given in tables 6, 7 and 8. Initial values for Moon's coefficients of potential are from LP150Q gravity model described in Konopliv (2001), and some of them are then fitted. Initial values for Earth's coefficients of potential are from EGM96 (<http://cddis.nasa.gov/926/egm96/egm96.html>); J_2 and J_3 are then fitted. Furthermore, two biases have been fitted:

- the first one on Cerga's observations between December 1996 and June 1998. It was first noticed by Chapront-Touzé et al. (2000). Its fitted value is -0.6 ns.
- the second one is applied on Mc Donald's observations, between October 1972 and August 1975. Its fitted value is 2.2 ns.

The final version of INPOP10a was obtained after an iterative process between planetary fit and Moon fit.

4 Results and Applications

The table 1 gives the planetary postfit residuals obtained with INPOP08 and INPOP10a. Some of the data sets used in the INPOP10a were not used in the INPOP08 adjustment: this explains some differences between the two columns of table 1. Postfit residuals are plotted in figures 3, 4, 5, and 6. The figures 9 show LLR residuals for each station and table 9 gives their standard deviations.

4.1 Asteroid masses

In table 2 and figures 1 and 2 are presented the asteroid masses obtained with INPOP10a compared to values found in the literature. The major sources of comparisons are the values obtained with DE421 (Folkner et al. 2009), DE423 (Konopliv et al. 2010) and Baer (2010) with realistic errorbars estimated by Kuchynka (2010). On plots 1 and 2, the comparisons of the asteroid GMs are ranked by their impact on the Earth-Mars distances over the 1990 to 2010 period. It then appears clearly that the estimations for the most perturbing objects are quite consistent when the estimations of the weak perturbing objects show bigger discrepancies. This result is consistent with the planetary ephemerides method for asteroid mass computation. For the asteroids inducing perturbations up to 10 meters, the differences in GMs are usually below 1-sigma or very close to 1-sigma except for 52 Europa. For this asteroid, which induced a maximum of 10 meters on the Earth-Mars distances, the errorbars are very large for all the determinations based on planetary ephemerides: we thus conclude to a bad determination of this mass based on the present interval of available data. In such case, one should have fixed this mass to the value estimated by other methods and especially by close-encounters as the one obtained by Baer et al (2008). This should be done for the next INPOP version. However, the good quality of the INPOP10a postfit and extrapolated residuals already tells us that such modifications would not change significantly the global behaviour of the ephemerides.

4.2 Extrapolation tests

Table 4 gives the residuals obtained by extrapolation of the two ephemerides INPOP08 and INPOP10a out of their fit interval. These residuals are plotted in figure 8. INPOP10a is fitted over the [1914.2:2009.7] time interval when INPOP08 was fitted over the interval [1914.2:2008.2]. For Venus, the extrapolation residuals are the same for INPOP08 and INPOP10a. For Mars, we first compare the extrapolation residuals of INPOP10a and of INPOP08 over the same time interval, from 2009.7 to 2010.35. Over this period, the residuals of INPOP10a are 9 times smaller than those obtained with INPOP08 (see table 4). This is obviously induced by the prolongation of the fitting interval and the addition of new data sets. In order to really test the improvement of the INPOP extrapolation capabilities, we compare the residuals obtained 6 months after the end of the fit period, on [2008.2:2008.8] for INPOP08 and over [2009.8:2010.2] for INPOP10a. Over these two intervals, the INPOP10a residuals have a clear linear drift of 20 meters/yr while the INPOP08 residuals show a more quadratic trend, as one can see in figure 9. This pure linear trend of INPOP10a extrapolation residuals can easily be removed (top right-hand plot of figure 9). This is an improvement compared to the residuals obtained with INPOP08. Indeed even after the correction of the linear trend, a quadratic signal is still remaining in the INPOP08 extrapolation residuals. This new behaviour of INPOP10a extrapolation capabilities could be induced by the changes in the fit of the asteroid masses.

4.3 Determination of sun J_2 and PPN β

Table 3 gives the obtained values for the mass of the Sun, the sun J_2 and the interval of sensitivity of data to modification of PPN β . Based on Fienga et al. (2010), we realized several fits for different values of β with a simultaneous fitting of initial conditions of planets, mass of the sun and asteroid densities. The given interval corresponds to values of β inducing changes in the postfit residuals below 5%. As one can see in table 3, we obtain with INPOP10a an improved estimation of β mainly due to the use in the fit of the Mariner and Messenger normal points of Mercury.

5 Work in progress

A complementary work is in progress to improve the physical interpretation of the fitted masses of asteroids. New estimations of supplementary advances of perihelia of planets (especially Saturn) will be computed rapidly as well as combined estimations of the PPN parameters β and γ .

6 References

Baer, J., 2010, <http://home.earthlink.net/~jimbaer1/astmass.txt>

Baer, J.; Milani, A.; Chesley, S.; Matson, R. D., 2008, *IAU Observational Error Model, and Application to Asteroid Mass Determination*, American Astronomical Society, DPS meeting 40, 52.09; Bulletin of the American Astronomical Society, Vol. 40, p.493.

Chapront-Touzé, M.; Chapront, J.; Francou, G., 2000, *Determination of UT0 with LLR observations*, Journées 1999 - systèmes de référence spatio-temporels, p. 217-220.

Fienga, A., ; Laskar, J.; Morley, T.; Manche, H.; Kuchynka, P.; Le Poncin-Lafitte, C.; Budnik, F.; Gastineau, M.; Somenzi, L., 2009, *INPOP08, a 4-D planetary ephemeris: from asteroid and time-scale computations to ESA Mars Express and Venus Express contributions*, Astronomy and Astrophysics, 507, 1675.

Fienga, A.; Laskar, J.; Kuchynka, P.; Le Poncin-Lafitte, C.; Manche, H.; Gastineau, M., 2010, *Gravity tests with INPOP planetary ephemerides*, Proceedings of the IAU Symposium "Relativity in Fundamental Astronomy: Dynamics, Reference Frames, and Data Analysis", Volume 261, p. 159-169.

Folkner, W.M., 2010, private communication.

Folkner, W.M., 2009, private communication.

Jones, Dayton L.; Fomalont, E.; Dhawan, V.; Romney, J.; Lanyi, G.; Border, J.; Folkner, W.; Jacobson, R., 2010, *Astrometric Observations of Cassini with the VLBA: The First Eight Epochs*, American Astronomical Society, AAS Meeting 215, 448.13; Bulletin of the American Astronomical Society, Vol. 42, p.456.

Konopliv et al., 2010, in press or submitted ?

Konopliv, A. S.; Asmar, S. W.; Carranza, E.; Sjogren, W. L.; Yuan, D. N., 2001, *Recent Gravity Models as a Result of the Lunar Prospector Mission*, Icarus, 150, p. 1-18

Kuchynka, P., Laskar, J., Fienga, A., Manche, H., 2010., *A ring as a model of main-belt in planetary ephemerides.*, Astronomy and Astrophysics, 514, A96.

Kuchynka, P., 2010, Paris Observatory PhD,

Lawson, C., Hanson, R., 1995, in *Solving Least Squares Problems, Revised edition*, eds. SIAM

Luzum, B.J., 2010, *System of Astronomical Constants*, Proceedings of the IAU Symposium "Relativity in Fundamental Astronomy: Dynamics, Reference Frames, and Data Analysis", Volume 261, p. 878.

Morley, T., 2010, private communication.

Morley, T., 2009, private communication.

Sicardy, B., 2009, private communication.

Table 1: Statistics of the residuals obtained after the INPOP10a fit. For comparisons, means and standard deviations of residuals obtained with INPOP08 are given.

Planet	Nbr	Time Interval	INPOP08	INPOP10a
Type of Data			mean $\pm 1\sigma$	mean $\pm 1\sigma$
Mercury				
Direct range [m]	462	1965-2000	30 $\pm$ 842	7 $\pm$ 866
Mariner range [m]	2	1974-1975	-1000 $\pm$ 305	-28 $\pm$ 85
Messenger Mercury flybys				
Messenger ra [mas]	3	2008-2009	1.1 $\pm$ 0.7	0.4 $\pm$ 1.2
Messenger de [mas]	3	2008-2009	2.0 $\pm$ 1.9	1.9 $\pm$ 2.1
Messenger range [m]	3	2008-2009	52 $\pm$ 619	-0.6 $\pm$ 1.9
Venus				
Direct range [km]	489	1965-2000	0.5 $\pm$ 2.3	0.5 $\pm$ 2.2
VEX range [m]	22145	2006-2010	1.6 $\pm$ 4.4	-0.2 $\pm$ 3.9
VLBI [mas]	22	1990-2007	2 $\pm$ 2	2 $\pm$ 2.5
Mars range [m]				
MGS	10474	1998-2008	-0.9 $\pm$ 1.6	0.5 $\pm$ 1.9
MEX	24262	2006-2010	-3.5 $\pm$ 2.0	0.0 $\pm$ 1.7
Path	90	1997	6.8 $\pm$ 12.5	-5.0 $\pm$ 5.0
Vkg	1256	1976-1982	-27.4 $\pm$ 19.0	-5.7 $\pm$ 35.0
Mars VLBI [mas]	96	1989-2007	0.5 $\pm$ 0.5	-0.0 $\pm$ 0.4
Jupiter flybys				
ra [mas]	5	1974-2000	48.0 $\pm$ 40.0	6 $\pm$ 5
de [mas]	5	1974-2000	-10.0 $\pm$ 50	-13 $\pm$ 18
range [km]	5	1974-2000	-27 $\pm$ 55	-0.6 $\pm$ 1.6
Jupiter VLBI [mas]	24	1996-1997	4 $\pm$ 11	0.2 $\pm$ 11
Jupiter Optical				
ra [mas]	6216	1914-2008	20 $\pm$ 304	-26 $\pm$ 304
de [mas]	6082	1914-2008	-44 $\pm$ 313	-54 $\pm$ 303
Saturn Cassini				
ra [mas]	31	2004-2007	1.5 $\pm$ 4	0.7 $\pm$ 4
de [mas]	31	2004-2007	7.0 $\pm$ 7	6.5 $\pm$ 7
range [m]	31	2004-2007	0.5 $\pm$ 22	0.0 $\pm$ 17
Saturn VLBI Cassini				
ra [mas]	10	2004-2009	0.3 $\pm$ 0.7	0.0 $\pm$ 0.6
de [mas]	10	2004-2009	-1.2 $\pm$ 2.0	0.1 $\pm$ 0.4
Saturn Optical				
ra [mas]	7824	1914-2008	-16 $\pm$ 305	-16 $\pm$ 305
de [mas]	7799	1914-2008	-7 $\pm$ 276	-9 $\pm$ 276
Uranus flybys				
ra [mas]	1	1986	-90	-30
de [mas]	1	1986	-36	-7
range [km]	1		1139	0.080
Uranus Optical				
ra [mas]	4145	1914-2008	-44 $\pm$ 278	-27 $\pm$ 290
de [mas]	4130	1914-2008	-38 $\pm$ 339	-11 $\pm$ 338
Neptune flybys				
ra [mas]	1	1989	-88	-11
de [mas]	1	1989	-48	-10
range [km]	1		2305	0.004
Neptune Optical				
ra [mas]	4340	1914-2008	-32 $\pm$ 282	2 $\pm$ 281
de [mas]	4320	1914-2008	-36 $\pm$ 335	2 $\pm$ 330
Pluto occultation				
ra [mas]	13	2005-2009	-6 $\pm$ 46	-1 $\pm$ 47
de [mas]	13	2005-2009	16 $\pm$ 30	-2 $\pm$ 19
Pluto Optical				
ra [mas]	2449	1914-2008	353 $\pm$ 926	38 $\pm$ 629
de [mas]	2463	1914-2008	-22 $\pm$ 524	17 $\pm$ 536

Table 2: Asteroids masses found in the recent literature and compared to the values estimated in INPOP10a. (F) is for fixed value.

Asteroids IAU	INPOP10a $10^{12} \times M_{\odot}$	Baer 2010 $10^{12} \times M_{\odot}$	DE421 $10^{12} \times M_{\odot}$	DE423 $10^{12} \times M_{\odot}$
1	475.836 ± 2.849	475.500 ± 4.755	468.488 ± 46.849	467.900 ± 3.250
2	111.394 ± 2.808	106.000 ± 1.060	100.979 ± 10.098	103.440 ± 2.550
4	133.137 ± 1.683	133.070 ± 0.266	132.835 ± 13.284	130.970 ± 2.060
7	7.772 ± 1.142	6.860 ± 1.029	5.998 ± 0.600	5.530 ± 1.320
324	4.692 ± 0.379		4.980 ± 0.498	5.340 ± 0.990
3	11.604 ± 1.313	13.400 ± 2.278	11.573 ± 1.157	12.100 ± 0.910
6	7.084 ± 1.212	6.460 ± 0.969	4.558 ± 0.456	6.730 ± 1.640
8	4.072 ± 0.631	4.260 ± 0.980	1.778 ± 0.178	2.010 ± 0.420
9	5.700 (F)	5.700 ± 1.425	4.272 ± 0.427	3.280 ± 1.080
10	44.500 (F)	44.500 ± 0.890	40.416 ± 4.042	44.970 ± 7.760
11	1.886 ± 1.029	3.090 ± 0.989	2.682 ± 0.268	
13	8.200 (F)	8.200 ± 1.640	3.104 ± 0.310	
14	4.130 (F)	4.130 ± 0.991	2.622 ± 0.262	1.910 ± 0.810
15	18.856 ± 1.617	15.700 ± 0.942	12.342 ± 1.234	14.180 ± 1.490
16	11.212 ± 5.174	11.000 ± 0.990	16.825 ± 1.682	12.410 ± 3.440
18	1.845 (F)		2.012 ± 0.201	
19	6.380 (F)	6.380 ± 1.021	3.489 ± 0.349	3.200 ± 0.530
20	2.850 (F)	2.850 ± 0.998	2.193 ± 0.219	
21	1.275 ± 1.170		1.047 ± 0.105	
24	2.826 ± 1.902	5.670 ± 2.155	3.036 ± 0.304	
25	0.002 ± 0.002		0.301 ± 0.030	
28	4.652 ± 0.990		1.243 ± 0.124	
29	5.920 (F)	5.920 ± 1.006	6.826 ± 0.683	7.420 ± 1.490
31	3.130 (F)	3.130 ± 1.002	8.582 ± 0.858	
41	9.213 ± 2.631		3.971 ± 0.397	4.240 ± 1.770
42	1.854 ± 1.070		0.693 ± 0.069	
45	2.860 (F)	2.860 ± 0.057	2.991 ± 0.299	
51	0.726 ± 0.421		1.085 ± 0.108	
52	42.304 ± 8.056	8.290 ± 0.995	10.202 ± 1.020	11.170 ± 8.400
60	0.402 ± 0.375		0.158 ± 0.016	
63	2.022 ± 1.685		0.769 ± 0.077	
65	7.173 ± 4.256	8.930 ± 0.982	5.229 ± 0.523	
78	3.232 ± 2.319		0.640 ± 0.064	
94	15.847 ± 11.522		3.119 ± 0.312	
105	1.109 (F)		0.663 ± 0.066	
107	18.205 ± 4.647	5.630 ± 0.169		
130	11.152 ± 8.032	3.320 ± 0.199		
135	0.917 ± 0.884		0.588 ± 0.059	
139	5.896 ± 3.315		1.417 ± 0.142	
145	2.266 (F)		1.138 ± 0.114	
187	2.484 ± 1.075		0.791 ± 0.079	
192	0.719 (F)		0.806 ± 0.081	
194	8.803 ± 2.921		1.371 ± 0.137	
216	0.564 ± 0.460		2.253 ± 0.225	
253	0.904 ± 0.650	0.052 ± 0.002		
337	0.543 ± 0.080		0.249 ± 0.025	
344	0.342 ± 0.188		0.859 ± 0.086	
354	2.451 (F)		2.464 ± 0.246	
372	4.443 (F)		2.675 ± 0.267	
419	0.997 ± 0.554		0.769 ± 0.077	
451	20.978 ± 14.797		4.596 ± 0.460	
488	6.234 ± 5.539		1.236 ± 0.124	
511	19.903 ± 4.068	22.000 ± 1.100	12.342 ± 1.234	8.580 ± 5.930
532	2.895 ± 0.759		6.676 ± 0.668	4.970 ± 2.810
554	1.575 ± 1.277		0.332 ± 0.033	
704	18.600 (F)	18.600 ± 1.116	18.565 ± 1.857	19.970 ± 6.570
747	6.032 ± 2.306		1.477 ± 0.148	
804	2.513 ± 1.820	2.020 ± 1.010		




Figure 1: Comparisons of the first 23 asteroid masses given in 10^{12} solar mass estimated by different authors and ranked by their impact on the Earth-Mars distances over 1990 to 2010. The mass estimations of the 3 biggest asteroids and perturbors Ceres, Pallas and Vesta are given in table 2. "others" stands for Baer (2010).




Figure 2: Comparisons of the other asteroid masses given in 10^{12} solar mass estimated by different authors and ranked by their impact on the Earth-Mars distances over 1990 to 2010. "others" stands for Baer (2010).

Table 3: Values of paramaters obtained by the fit of INPOP08 and INPOP10a to observations. In INPOP08, the mass of the sun was not fitted directly but deduced from the fitted value of AU (Fienga et al. 2009). The (F) indicates that the marked values are fixed in the fit. The published uncertainty (Standish 1998) of the INPOP08 $GM_{\odot}$ fixed value is $\pm 50 \text{ km}^3. \text{ s}^{-2}$. The equivalent value of AU deduced from the estimation of the $GM_{\odot}$ in INPOP10a is given in line 6. The INPOP σ for the Earth-Moon mass ratio EMRAT, the $J2_{\odot}$ and the PPN parameter β are the limit of sensitivity of the observations to the tested parameter. For the AU and the $GM_{\odot}$, the σ are deduced from the least squares fit.

	INPOP08 $\pm 1\sigma$	INPOP10a $\pm 1\sigma$	DE423
EMRAT	(81.30054 ± 0.00005)	$(81.3005700 \pm 0.0000010)$	$(81.3005694 \pm 0.0000015)$
$J2_{\odot}$	$(1.82 \pm 0.47) \times 10^{-7}$	$(2.40 \pm 0.25) \times 10^{-7}$	
$(\beta - 1) \times 10^{-4}$	(0.75 ± 1.25)	(0.25 ± 0.75)	(0.4 ± 2.4)
$GM_{\odot} [\text{km}^3. \text{ s}^{-2}]$	132712440017.98700 (F)	132712440055 ± 1	132712440042 ± 10
AU [m]	149597870699.2 ± 0.11	149597870691.0 (F)	
AU [m] from $GM_{\odot}$		149597870704.9 ± 0.3	149597870700.0 ± 3

Table 4: Residuals obtained by extrapolation of INPOP08 and INPOP10a out from the fit interval.

Planet Type of Data		INPOP08 mean $\pm 1\sigma$	INPOP10a mean $\pm 1\sigma$
Mars range [m]			
MEX	2009.7:2010.35	-18 ± 63	11.3 ± 8.6
VEX	2009.7:2010.35	17 ± 50	18 ± 50

Table 5: Masses of asteroids fitted during the fit of INPOP10a. The σ is deduced from the least squares fit.

Asteroids	M	σ	Asteroids	M	σ	Asteroids	M	σ
IAU	$10^{12} \times M_{\odot}$	$10^{12} \times M_{\odot}$		$10^{12} \times M_{\odot}$	$10^{12} \times M_{\odot}$		$10^{12} \times M_{\odot}$	$10^{12} \times M_{\odot}$
1	475.836	2.849	84	0.275	0.204	381	4.615	0.400
2	111.394	2.808	89	4.181	2.083	386	11.825	0.210
4	133.137	1.683	94	15.847	11.522	387	2.674	0.310
7	7.772	1.142	105	1.109	(F)	404	1.718	1.522
324	4.692	0.379	107	18.205	4.647	410	3.476	2.380
3	11.604	1.313	112	0.990	3.410	416	1.643	1.560
6	7.084	1.212	117	8.659	0.100	419	0.997	0.554
8	4.072	0.631	126	0.237	2.910	420	7.417	0.460
9	5.700	(F)	127	4.402	0.440	442	0.098	0.010
10	44.500	(F)	128	4.583	2.772	444	7.992	5.581
11	1.886	1.029	129	4.336	1.846	445	1.744	0.390
12	3.774	1.679	130	11.152	8.032	449	0.791	0.703
13	8.200	(F)	132	0.207	1.360	451	20.978	14.797
14	4.130	(F)	135	0.917	0.884	455	0.596	0.060
15	18.856	1.617	138	0.248	0.130	469	3.647	3.140
16	11.212	5.174	139	5.896	3.315	471	6.359	0.360
18	1.845	(F)	141	4.146	2.902	481	2.908	0.730
19	6.380	(F)	144	4.566	2.977	485	0.686	0.220
20	2.850	(F)	145	2.266	(F)	488	6.234	5.539
21	1.275	1.170	147	6.185	0.240	491	2.425	0.980
24	2.826	1.902	148	2.459	0.840	503	1.434	0.170
25	0.002	0.002	150	9.085	0.420	505	2.008	1.931
26	0.376	0.450	156	3.263	1.864	511	19.903	4.068
28	4.652	0.990	163	1.008	0.340	516	0.720	0.668
29	5.920	(F)	164	0.467	0.390	532	2.895	0.759
31	3.130	(F)	165	9.761	0.100	554	1.575	1.277
33	3.117	0.370	168	8.599	0.950	582	0.215	0.590
34	1.816	1.703	173	6.743	5.668	584	0.414	0.290
36	2.170	1.909	187	2.484	1.075	602	5.107	0.240
38	2.872	2.748	192	0.719	(F)	604	0.728	0.140
39	8.799	0.510	194	8.803	2.921	626	2.691	3.280
41	9.213	2.631	200	5.387	0.810	665	0.351	0.200
42	1.854	1.070	204	0.302	0.910	675	6.058	1.200
43	0.753	0.430	210	1.713	0.550	679	0.359	0.100
45	2.860	(F)	211	3.801	0.660	680	1.352	0.020
46	3.525	2.630	212	6.639	0.530	690	6.428	0.140
47	5.387	0.230	216	0.564	0.460	702	6.859	5.606
48	20.106	14.987	217	0.765	0.030	704	18.600	(F)
50	1.833	1.344	221	2.951	0.170	735	1.081	0.340
51	0.726	0.421	234	0.221	0.730	739	0.584	0.540
52	42.304	8.056	240	2.067	1.912	747	6.032	2.306
53	2.830	2.515	253	0.904	0.650	751	3.552	0.260
54	2.237	1.113	266	2.085	0.210	758	0.468	0.040
56	2.321	1.812	268	0.786	0.680	760	0.668	0.662
59	2.587	1.795	304	0.576	0.563	769	3.174	0.320
60	0.402	0.375	306	0.268	0.290	784	1.882	0.160
61	1.454	1.400	322	0.936	0.020	786	1.416	1.401
63	2.022	1.685	328	4.889	0.130	804	2.513	1.820
65	7.173	4.256	337	0.543	0.080	914	1.183	0.120
67	0.516	0.050	344	0.342	0.188	949	0.872	0.310
72	1.668	4.270	345	2.195	0.230	1013	0.086	0.720
74	3.082	2.696	346	3.182	0.090	1015	2.398	0.340
77	0.874	0.340	354	2.451	(F)	1021	2.585	0.060
78	3.232	2.319	365	2.937	0.480	1036	0.084	0.160
81	3.111	2.671	372	4.443	(F)	1171	0.908	0.100

Table 6: Values of dynamical parameters fitted to LLR observations. GM_{EMB} is the sum of Earth's and Moon's masses, multiplied by the gravitationnal constant and is expressed in AU^3/day^2 . C_{nmE} are the Earth's coefficients of potential (without unit). τ_{21E} and τ_{22E} are time delays of the Earth used for tides effects and expressed in days. C_{nmM} and S_{nmM} are the Moon's coefficients of potential (without unit). $(C/MR^2)_M$ is the ratio between the third moment of inertia of the Moon, divided by its mass and the square of the mean equatorial radius (without unit). k_{2M} and τ_M are Love number (without unit) and time delay (in day) of the Moon. Formal errors at 1σ are given if the parameter is fitted. Fixed values come from Lunar gravity model LP15Q or Earth's one EGM96.

Name	Value	Formal error (1σ)
GM_{EMB}	$8.9970116036 \times 10^{-10}$	$\pm 7.2 \times 10^{-19}$
C_{20E}	$-1.0826282367 \times 10^{-3}$	$\pm 1.1 \times 10^{-9}$
C_{30E}	$2.8862851809 \times 10^{-6}$	$\pm 3.9 \times 10^{-8}$
C_{40E}	$1.6196215914 \times 10^{-6}$	
τ_{21E}	$1.2326471124 \times 10^{-2}$	$\pm 9.7 \times 10^{-5}$
τ_{22E}	$6.9795966104 \times 10^{-3}$	$\pm 7.7 \times 10^{-6}$
C_{20M}	$-2.0347464196 \times 10^{-4}$	$\pm 2.8 \times 10^{-8}$
C_{22M}	$2.2400822245 \times 10^{-5}$	$\pm 2.7 \times 10^{-9}$
C_{30M}	$-8.3743483591 \times 10^{-6}$	$\pm 2.3 \times 10^{-8}$
C_{31M}	$3.1934585938 \times 10^{-5}$	$\pm 3.7 \times 10^{-7}$
C_{32M}	$4.8452131770 \times 10^{-6}$	
C_{33M}	$1.7262951856 \times 10^{-6}$	$\pm 6.3 \times 10^{-9}$
C_{40M}	$9.6422863508 \times 10^{-6}$	
C_{41M}	$-5.6926874003 \times 10^{-6}$	
C_{42M}	$-1.5861997683 \times 10^{-6}$	
C_{43M}	$-8.1204110561 \times 10^{-8}$	
C_{44M}	$-1.2739414703 \times 10^{-7}$	
S_{31M}	$3.2781092481 \times 10^{-6}$	$\pm 8.8 \times 10^{-8}$
S_{32M}	$1.6873163963 \times 10^{-6}$	$\pm 5.7 \times 10^{-10}$
S_{33M}	$-2.4855254932 \times 10^{-7}$	
S_{41M}	$1.5743934837 \times 10^{-6}$	
S_{42M}	$-1.5173124037 \times 10^{-6}$	
S_{43M}	$-8.0279066453 \times 10^{-7}$	
S_{44M}	$8.3147478750 \times 10^{-8}$	
$(C/MR^2)_M$	$3.9319024676 \times 10^{-1}$	$\pm 4.7 \times 10^{-5}$
k_{2M}	$2.6248414243 \times 10^{-2}$	$\pm 1.7 \times 10^{-4}$
τ_M	$1.8896343615 \times 10^{-1}$	$\pm 1.2 \times 10^{-3}$

Table 7: Selenocentric coordinates of reflectors, expressed in meters.

Reflector		Value	Formal error (1σ)
Apollo 11	x	1591926.238	± 1.510
	y	690799.993	± 3.460
	z	21002.457	± 0.306
Apollo 14	x	1652725.404	± 1.140
	y	-520893.461	± 3.600
	z	-109731.862	± 0.318
Apollo 15	x	1554675.870	± 0.276
	y	98193.648	± 3.380
	z	765004.763	± 0.304
Lunakhod2	x	1339316.595	± 1.750
	y	801956.645	± 2.910
	z	756358.336	± 0.267

Table 8: ITRF coordinates of stations at J1997.0, expressed in meters.

Station		Value	Formal error (1σ)
Cerga	x	4581692.121	± 0.003
	y	556196.025	± 0.001
	z	4389355.020	± 0.010
Mc Donald	x	-1330781.440	± 0.012
	y	-5328755.514	± 0.009
	z	3235697.496	± 0.022
MLRS1	x	-1330121.099	± 0.014
	y	-5328532.296	± 0.008
	z	3236146.576	± 0.024
MLRS2	x	-1330021.431	± 0.002
	y	-5328403.289	± 0.003
	z	3236481.603	± 0.010
Haleakala (rec.)	x	-5466000.459	± 0.011
	y	-2404424.720	± 0.013
	z	2242206.708	± 0.028
Apollo	x	-1463998.834	± 0.005
	y	-5166632.675	± 0.004
	z	3435013.101	± 0.012

Table 9: Means and standard deviations of LLR residuals for INPOP10a solution; both are expressed in centimeters. Na is the total number of observations available, Nk is the number kept in fitting process, Nr is the number that have been rejected according to the 3σ criterion ($Na=Nk+Nr$).

Station	Period	Mean	Std. dev.	Na	Nk	Nr
Cerga	1987-1995	-0.16	6.37	3460	3415	45
Cerga	1995-2010	-0.03	4.00	4932	4861	71
Cerga	1984-1986	8.11	16.00	1187	1158	29
Mc Donald	1969-1986	0.16	31.78	3604	3489	115
MLRS1	1982-1985	-7.89	73.28	418	405	13
MLRS1	1985-1988	0.22	7.30	174	163	11
MLRS2	1988-1996	-0.85	4.27	1192	1148	44
MLRS2	1996-2008	0.56	4.82	2243	1768	475
Haleakala	1984-1990	-0.44	8.10	770	734	36
Apollo	2006-2009	0.08	4.89	642	640	2




Figure 3: Postfit residuals obtained with INPOP10a




Figure 4: Postfit residuals obtained with INPOP10a




Figure 5: Postfit residuals obtained with INPOP10a




Figure 6: Postfit residuals obtained with INPOP10a




Figure 7: Extrapolation of MEX and VEX out from the fit interval.




Figure 8: Zooms from Figure 8 plotting the MEX extrapolated residuals.




Figure 9: Postfit LLR residuals with INPOP10a for each station, expressed in centimeters.

Asteroids IAU	INPOP10a $10^{12} \times M_{\odot}$	Baer 2010 $10^{12} \times M_{\odot}$	DE421 $10^{12} \times M_{\odot}$	DE423 $10^{12} \times M_{\odot}$
1	475.836 ± 2.849	475.500 ± 4.755	468.488 ± 46.849	467.900 ± 3.250
2	111.394 ± 2.808	106.000 ± 1.060	100.979 ± 10.098	103.440 ± 2.550
4	133.137 ± 1.683	133.070 ± 0.266	132.835 ± 13.284	130.970 ± 2.060

	INPOP08 $\pm 1\sigma$	INPOP10a $\pm 1\sigma$	DE423 $\pm 1\sigma$
EMRAT	(81.30054 ± 0.00005)	$(81.3005700 \pm 0.0000010)$	$(81.3005694 \pm 0.0000015)$
$J2_{\odot}$	$(1.82 \pm 0.47) \times 10^{-7}$	$(2.40 \pm 0.25) \times 10^{-7}$	
$(\beta - 1) \times 10^{-4}$	(0.75 ± 1.25)	(0.25 ± 0.75)	(0.4 ± 2.4)
	$\pm 5\sigma$	$\pm 5\sigma$	$\pm 1\sigma$
$GM_{\odot} [\text{km}^3 \cdot \text{s}^{-2}]$	132712440017.98700 (F)	132712440055 ± 1	132712440042 ± 10
AU [m]	149597870699.2 ± 0.55	149597870691.0 (F)	
AU [m] from $GM_{\odot}$		149597870704.9 ± 0.3	149597870700.0 ± 3
	$\pm 1\sigma$	$\pm 1\sigma$	$\pm 1\sigma$
Ceres [$10^{12} \times M_{\odot}$]	465.8 ± 4.5	475.836 ± 2.849	467.900 ± 3.250
Pallas [$10^{12} \times M_{\odot}$]	107.6 ± 10.0	111.394 ± 2.808	103.440 ± 2.550
Vesta [$10^{12} \times M_{\odot}$]	139.2 ± 15.0	133.137 ± 1.683	130.970 ± 2.060

Table 1: Statistics of the residuals obtained after the INPOP10a fit. For comparisons, means and standard deviations of residuals obtained with INPOP08 are given.

Planet Type of Data	Nbr	Time Interval	INPOP08 mean $\pm 1\sigma$	INPOP10a mean $\pm 1\sigma$
Mercury				
Direct range [m]	462	1965-2000	30 ± 842	7 ± 866
Mariner range [m]	2	1974-1975	-1000 ± 305	-28 ± 85
Messenger Mercury flybys				
Messenger ra [mas]	3	2008-2009	1.1 ± 0.7	0.4 ± 1.2
Messenger de [mas]	3	2008-2009	2.0 ± 1.9	1.9 ± 2.1
Messenger range [m]	3	2008-2009	52 ± 619	-0.6 ± 1.9